arXiv:1407.8473v1 [math-ph] 31 Jul 2014

A.Cormack's last inversion formula and a FBP reconstruction

V. P. Palamodov

28.07.2014

Abstract

A reconstruction of a function from integrals over the family of confocal paraboloids in $\mathbb{R}^n$ is given by a FBP formula.

1 Introduction

Cormack proposed a clever method for determination of distribution of a scattered material in a space from data of scattered light filtered on travel times [5]. He wrote in the introduction to [5]: "Suppose that waves travel in this space with a speed v, and that the space contains a distribution of scattering material with a density f which is assumed to be smooth and either rapidly decreasing or of compact support... An impulsive plane wave front with a normal ω reaches 0 at $t = 0$. As result of the well-known focusing property of paraboloids the scattering waves arriving at 0 at time $t = 2p/v$ will all have originated on the paraboloid defined by $r\,(1 + \langle \xi, \omega \rangle) = 2p$..." The mathematical problem is to reconstruct a function in a plane from data of integrals $\mathrm{R}f$ over a family of confocal parabolas or confocal rotation paraboloids. "The problem of determining f from $\mathrm{R}f$ for a family of curves in $\mathbb{R}^2$ was discussed in [1], [2], [3] and in [3] it was shown that the parabolic case could be reduced to the ordinary Radon transform by a coordinate transformation which does not have an analog in $\mathbb{R}^3$. The case of $\mathbb{R}^n$ was considered in [4], in which f was considered to be expanded in spherical harmonics; some properties of the harmonic components were given, and a solution to the resulting integral equation was given. This solution, however, had a very awkward form and as a result it was not investigated in detail. Given in [5] is a simpler solution for the case of $\mathbb{R}^3$..." This solution is

$$r^{1/2} f\,(x) = -\frac{1}{(4\pi)^2} \int_{\mathrm{S}^2} \left(\frac{\partial}{\partial r} r\right)^2 \frac{g\,(p,\omega)}{p^{3/2}}\bigg|_{p=r(1+\langle \xi,\omega\rangle)/2} \Omega, \tag{1}$$

where $r = |x|\,,\ \xi = x/\,|x|$, Ω is the area form in S^2 and g is the surface integral of f.

Our goal is a reconstruction by means of an integral transform of FBP type to avoid divergent integrals.

2 Confocal paraboloids and integrals

Confocal parabolas



Reflecting paraboloid



The family of paraboloids $Z(p,\omega)$ in an Euclidean space E^n with the focus at the origin can be given by the equation $r-\langle x,\omega\rangle = 2p$ where $p>0$ and $\omega \in \mathrm{S}^{n-1}$ are parameters. For a bounded function f in E^n with compact support we consider the integrals over paraboloids

$$\mathrm{R}f(p,\omega) = \int_{Z(p,\omega)} f\mathrm{d}S,$$
$$\mathrm{M}f(p,\omega) = \lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\int_{p\le r-\langle x,\omega\rangle\le p+\varepsilon} f\mathrm{d}x = \int_{Z(p,\omega)}\frac{f\mathrm{d}S}{|\nabla\theta(x,\omega)|}, \tag{2}$$

where $\mathrm{d}S$ is the Euclidean surface density and $\mathrm{M}f$ is equal to the integral of $f\mathrm{d}x$ between to close paraboloids. The function can be reconstructed from $\mathrm{R}f$ and from $\mathrm{M}f$ according to the following

Theorem 1 *For an arbitrary function $f \in C^2\left(E^2\right)$ with compact support, a reconstruction is given by*

$$f(x) = -\frac{1}{4\pi^2 r^{1/2}} \int_{\mathrm{S}^1} \int_{\mathbb{R}} \frac{\partial}{\partial p} \left(p^{1/2} \mathrm{R} f(p,\omega) \right) \frac{\mathrm{d}p}{\theta(x,\omega) - p} \Omega \tag{3}$$
$$= -\frac{1}{4\pi^2 r} \int_{\mathrm{S}^1} \int_{\mathbb{R}} \frac{\partial}{\partial p} \left(p \mathrm{M} f(p,\omega) \right) \frac{\mathrm{d}p}{\theta(x,\omega) - p} \Omega,\ x \in E^2 \backslash 0.$$

For $f \in C^2\left(E^3\right)$, we have

$$f(x) = -\frac{1}{8\pi^2 r^{1/2}} \int_{\mathrm{S}^2} \frac{\partial^2}{\partial p^2}\ p^{1/2} \mathrm{R} f(p,\omega) \Big|_{p=\theta(x,\omega)} \Omega \tag{4}$$
$$= -\frac{1}{8\pi^2 r} \int_{\mathrm{S}^2} \frac{\partial^2}{\partial p^2}\ p \mathrm{M} f(p,\omega)|_{p=\theta(x,\omega)} \Omega,\ x \in E^3 \backslash 0.$$

Remark. The first formula (4) reminds Cormack's reconstruction (8) where apparently $g = \mathrm{R}f$. However the integral $\mathrm{M}f$ is a correct model for the "scattered light filtrated on travel times". It is different from the Euclidean surface integral $\mathrm{R}f$ since the dominator in (2) is not a constant.

Proof. We assume at the beginning that $n \geq 2$ is arbitrary. The family of rotation paraboloids is generated by the function $\Phi(x,p,\omega) = \theta(x,\omega) - p$, $\theta(x,\omega) = (r - \langle x,\omega\rangle)/2$ defined in the manifold $X \times \Sigma$, $X = E^n \backslash 0$, $\Sigma = \mathbb{R}_+ \times \mathrm{S}^{n-1}$. The set X is supplied with the Euclidean volume form $\mathrm{d}x$ and the manifold Σ with the volume form $\mathrm{d}p \wedge \Omega$. The map $\pi_X : Z \to X$ is proper where $Z = \Phi^{-1}(0)$ and the function Φ but satisfies the conditions: π_X has rank n and the map π^* is a bijection see §3. We have for any $n \geq 2$

$$\nabla_x \theta = \frac{\xi - \omega}{2}, \text{ where } \xi = \frac{x}{r}, \text{ and } |\nabla_x \theta| = \theta^{1/2} r^{-1/2},$$
$$2^n (n-1)! J(x,\omega) = \nabla\theta \wedge \left(\mathrm{d}_\omega \nabla\theta\right)^{\wedge n-1} = \langle \xi - \omega, \omega\rangle\ \ \omega \wedge \left(\mathrm{d}\omega\right)^{\wedge n-1} = \langle \xi - \omega, \omega\rangle\, \Omega.$$

It follows that $\det J(x,\omega) \neq 0$ except for the submanifold $V \doteqdot \{\omega = \xi\}$ in $X \times \Sigma$. However π^* is a bijection since $\langle \xi - \omega, \omega\rangle \geq 0$. Choose a weight function $b(x,\omega) = |\nabla_x \theta(x,\omega)|^2$ and consider the weighted Funk transform

$$\mathrm{M}_b f(p,\omega) = \int_{\theta(x,\omega)=p} f(x) |\nabla_x \theta(x,\omega)| \, \mathrm{d}S = p^{1/2} \mathrm{R}\left(r^{-1/2} f\right) = p\mathrm{M}\left(r^{-1} f\right), \tag{5}$$

where R is the Euclidean integral transform as above. We want to invert M_b by means of a filtered back projection. Take a number $\varepsilon > 0$ and consider the ε-neighborhood $V(\varepsilon)$. The generating function Φ satisfies (**i**) for the map π^* restricted to the set $Z \backslash V(\varepsilon) \times \mathbb{R}_+$. Let $b(\varepsilon)$ be a function in $X \times \Sigma$ that vanishes in $V(\varepsilon)$ and coincides with b otherwise. Take $\beta = 1$ as the second weight functions and apply Theorem 5§3 to Φ and $Z \backslash V(\varepsilon)$. Note that equation $\mathrm{d}_\omega \theta(x,\omega) = -x/2$ implies (**ii**). This yields

$$D_{b(\varepsilon)} f = \mathrm{N}\mathrm{M}_{b(\varepsilon)} f + \Theta_{b(\varepsilon)} f,$$

where $\mathrm{N} = \mathrm{N}_\beta$ and $\Theta_{b(\varepsilon)}$ is the operator with the kernel

$$\Theta_{b(\varepsilon)}(x,y) = \mathrm{Re}\, i^n \int_{\mathrm{S}^2} \frac{b(\varepsilon)(y,\omega)\,\Omega}{(\theta(x,\omega) - \theta(y,\omega) - i0)^n}.$$

By (11)

$$D_{b(\varepsilon)}(x)=\frac{1}{|\mathrm{S}^{n-1}|}\int_{\mathrm{S}^{n-1}}\frac{b(\varepsilon)\,\Omega}{|\nabla_x\theta|^n}. \tag{6}$$

Lemma 2 *We have* $D_{b(\varepsilon)}(x)\to D_b(x)$ *for* $x\neq 0$ *and* $\Theta_{b(\varepsilon)}(x,y)\to 0$ *for arbitrary* $x\neq y\in E^n\backslash 0$ *as* $\varepsilon\to 0$ *where*

$$D_b(x)=\frac{1}{|\mathrm{S}^{n-1}|}\int_{\mathrm{S}^{n-1}}\frac{b\Omega}{|\nabla_x\theta|^n}=\frac{1}{|\mathrm{S}^{n-1}|}\int_{\mathrm{S}^{n-1}}\frac{\Omega}{|\nabla_x\theta|^{n-2}}=2^{n-2} \tag{7}$$

for any $n\geq 2$.

Proof of Lemma 2. We have $|\nabla_x\theta|^2=r^{-1}\theta$, hence $|\nabla\theta(x,\omega)|^{-1}\approx\sin\gamma/2$ where γ is the angle between x and ω. Therefore integral (7) converges to (6). This proves the first statement. To check the second statement we note that

$$\theta(x,\omega)-\theta(y,\omega)=\frac{1}{2}\left(|x|-|y|-\langle x-y,\omega\rangle\right)$$

is a linear function of ω for arbitrary $x\neq y$. This function has a zero $\omega\in\mathrm{S}^{n-1}$ since $||x|-|y||\leq|x-y|$. By [6] it follows that $\Theta_b(x,y)=0$ since $\theta(x,\omega)-\theta(y,\omega)$ is a linear function of ω and $\deg b+n-1=n$. Now it is sufficient to check that $\Theta_{b(\varepsilon)}\to\Theta_b$ as $\varepsilon\to 0$. For this we show that the dominator $\theta(x,\omega)-\theta(y,\omega)$ does not vanish at the point $\omega=\xi\doteq x/r$. We have

$$\theta(x,\xi)-\theta(y,\xi)=\frac{-|x|\,|y|+\langle y,x\rangle}{2|x|}<0$$

since $y\neq x$ and $x\neq 0, y\neq 0$. This completes the proof. ▶

By Lemma 2 we can take the limit in (10) and obtain the equation $D_bf=\mathrm{NM}_bf$. Due to (5) we can replace the integrals M_bf by data of $\mathrm{M}\left(r^{-1}f\right)$ and or by data of $\mathrm{R}\left(r^{-1/2}f\right)$. By substitution $f=r\tilde{f}$ or $f=r^{1/2}\tilde{f}$ we finally come up to the equations (3) and (4). This completes the proof of Theorem 1. ▶

Proposition 3 *Integrals (3) and (4) have the classical sense for sufficiently smooth functions* f.

Proof. For $n=2$, we have

$$\mathrm{d}Z=\left(\mathrm{d}r^2+r^2\mathrm{d}\gamma^2\right)^{1/2}=\frac{\mathrm{d}r}{\cos\gamma/2}=\frac{\mathrm{d}r}{(1-p/r)^{1/2}}$$

and

$$\begin{aligned}p^{1/2}\mathrm{R}\left(r^{1/2}f\right)(p,\omega)&=p^{1/2}\int_{\sin^2(\gamma/2)=p/r}r^{1/2}f(r,\gamma)\,\mathrm{d}Z\\&=2\int_p^\infty f(r,\gamma)\frac{r\mathrm{d}r}{(r/p-1)^{1/2}}=2\int_1^\infty f(sp,\gamma)\frac{s\mathrm{d}s}{(s-1)^{1/2}}.\end{aligned}$$

The right hand side has continuous second derivative with respect to p and ω if $f\in C^2\left(E^2\right)$.

Lemma 4 *For any* $f\in C^4\left(E^3\right)$, *we have*

$$\mathrm{M}f(p,\omega)=p^{-1/2}\mathrm{R}\left(r^{1/2}f\right)(p,\omega)=g_0(\omega)+pg_1(p,\omega)$$

as $p\to 0$ *where* g_0 *and* g_1 *are* C^2*-continuous functions.*

Theorem 1 will follow since $\mathrm{M}f = p^{-1/2}\mathrm{R}\left(r^{1/2}f\right)$ and

$$\frac{\partial^2}{\partial p^2}\left(p\mathrm{M}f\left(p,\omega\right)\right) = \frac{\partial^2}{\partial p^2}\left(p^{1/2}\mathrm{R}\left(r^{1/2}f\right)\left(p,\omega\right)\right) = 2g_1\left(p,\omega\right) + o\left(1\right).$$

Proof. Variables $r,\ \gamma$ and φ are spherical coordinates in $\mathbb{R}^3$ where $\gamma,\ 0 \leq \gamma < \pi$ is the spherical distance between points $\xi, \omega \in \mathrm{S}^2$ and φ is rotation angle about ω and $Z\left(p,\omega\right)$. Equation $p = \left(x - \langle x,\omega\rangle\right)/2 = r\sin^2\left(\gamma/2\right)$ defines the surface $Z\left(p,\omega\right)$ and we have

$$\mathrm{d}Z = \left(\mathrm{d}r^2 + r^2\mathrm{d}\gamma^2\right)^{1/2} r\sin\gamma\mathrm{d}\varphi$$

for the Euclidean area $\mathrm{d}Z$ of $Z\left(p,\omega\right)$. We have

$$\left(\mathrm{d}r^2 + r^2\mathrm{d}\gamma^2\right)^{1/2} = \frac{\mathrm{d}r}{\cos\gamma/2},\ \mathrm{d}Z = 2\left(pr\right)^{1/2}\mathrm{d}r\mathrm{d}\varphi.$$

This yields

$$p^{-1/2}\mathrm{R}\left(r^{1/2}f\right)\left(p,\omega\right) = p^{-1/2}\int_{\sin^2(\gamma/2)=p/r} r^{1/2}f\left(r,\gamma,\varphi\right)\mathrm{d}Z = 2\int_p^\infty\int_0^{2\pi} rf\left(r,\gamma,\varphi\right)\mathrm{d}\varphi\mathrm{d}r.$$

For a plane P and a function $\phi \in C^4\left(P\right)$, the circle integral can be represented in the form

$$\int_{|y|=s}\phi\left(y\right)\mathrm{d}\varphi = 2\pi\phi\left(0\right) + \tau\left(s\right)s^2$$

with a remainder $\tau \in C^2$. Applying this equation to f in a plane P orthogonal to ω for $s = r\sin\gamma$ we obtain

$$\int_0^{2\pi} f\left(r,\gamma,\varphi\right)\mathrm{d}\varphi = 2\pi f\left(r,0.\varphi\right) + \left(r\sin\gamma\right)^2 h\left(r,\gamma,\varphi\right),\ h \in C^2.$$

We have $\left(r\sin\gamma\right)^2 = 4p\left(r-p\right)$ and $f\left(r,0,\varphi\right) = f\left(r\omega\right)$ which implies

$$p^{-1/2}\mathrm{R}\left(r^{1/2}f\right)\left(p,\omega\right) = 4\pi\int_p^\infty rf\left(r\omega\right)\mathrm{d}r + 8p\int_p^\infty \tau\left(r,\gamma,\omega\right)\left(r-p\right)r\mathrm{d}r.$$

Now Lemma 4 follows for the functions

$$g_0\left(\omega\right) = 4\pi\int_p^\infty rf\left(r\omega\right)\mathrm{d}r,\ g_1\left(p,\omega\right) = 8\int_p^\infty \tau\left(r,\gamma,\omega\right)\left(r-p\right)r\mathrm{d}r$$

which belong to C^2. ▶

3 Weighted integral transform and reconstruction

Let $X,\ \Sigma$ be smooth manifolds of dimension $n > 1$ and $Z \subset X \times \Sigma$ be a hypersurface. Let $\pi_X : Z \to X,\ \pi_\Sigma : Z \to \Sigma$ be the natural projections. A family of hypersurfaces $Z\left(\sigma\right) = \pi_\Sigma^{-1}\left(\sigma\right)$ in X is defined as well as the family of hypersurfaces $Z\left(x\right) = \pi_X^{-1}\left(x\right)$ in Σ. Suppose that there exists a real function $\Phi \in C^2\left(X \times \Sigma\right)$ (called generating function) such that $Z = \Phi^{-1}\left(0\right)$ and $\mathrm{d}\Phi \neq 0$ in Z. Let $T^*\left(X\right)$ be the cotangent bundle of X and $T^*\left(X\right)/\mathbb{R}_+$ be the bundle of rays in $T^*\left(X\right)$. We suppose that

(**i**) The map $\pi^* : Z\times\mathbb{R}_+ \to T^*(X)$ is a diffeomorphism, where $\pi^*(x,\sigma;t) = (x, t\mathrm{d}_x\Phi(x,\sigma))$, $t \in \mathbb{R}_+$. This condition implies that π_X has rank n and $Z(\sigma)$ is a smooth hypersurface in X for any $\sigma \in \Sigma$. The function $\det J(\Phi)$ vanishes if where Π^* is not a local diffeomorphism and vice versa.

Definition. We say that points $x \neq y \in X$ are conjugate with respect to Φ, if $\Phi(x,\sigma) = \Phi(y,\sigma)$ and $\mathrm{d}_\sigma\Phi(x,\sigma) = \mathrm{d}_\sigma\Phi(y,\sigma)$ for some $\sigma \in \Sigma$. We call a generating function Φ *regular,* if the projection $\pi : Z \to X$ is proper, Φ satisfies the conditions (**i**) and (**ii**): there are no conjugate points.

Let $\mathrm{d}X$ be a volume form in X and $b = b(x,\sigma)$ be a continuous function in Z. For any bounded arbitrary function f in X with compact support, the integral

$$\mathrm{M}_b f(\sigma) \doteqdot \int \delta(\Phi(x,\sigma)) f(x) b(x,\sigma)\,\mathrm{d}X = \lim_{\varepsilon\to 0}\frac{1}{2\varepsilon}\int_{|\Phi(\cdot,\sigma)|\leq\varepsilon} f(x) b(x,\sigma)\,\mathrm{d}X$$

converges for any $\sigma \in \Sigma$. We suppose now that $\mathrm{d}X$ is the Riemannian volume form of a Riemannian metric g in X. We have:

$$\mathrm{M}_b f(\sigma) = \int_{Z(\sigma)} \frac{f(x) b(x,\sigma)}{|\mathrm{d}_x\Phi(x,\sigma)|_{\mathrm{g}}}\mathrm{d}_{\mathrm{g}}S, \tag{8}$$

where $\mathrm{d}_{\mathrm{g}}S$ is the Riemannian volume form in a hypersurface in X. Here $Z(\sigma) = \{x \in X, \Phi(x,\sigma) = 0\}$ and $|\cdot|_{\mathrm{g}}$ means the Riemannian norm of a covector. Let β be a locally bounded function in $X \times \Sigma$. The singular integral

$$\Theta_{b,\beta}(x,y) = (-1)^n \int_{Z(y)} \frac{b(y,\sigma)\beta(x,\sigma)}{(\Phi(x,\sigma) - i0)^n}\frac{\mathrm{d}\Sigma}{\mathrm{d}_\sigma\Phi(y,\sigma)} \tag{9}$$

is well defined for any $x, y \in X$, $y \neq x$ since $\mathrm{d}_\sigma\Phi(x,\sigma) \neq 0$ in $Z(y) \doteqdot \{\sigma \in \Sigma; \Phi(y,\sigma) = 0\}$ due to (**ii**).

Theorem 5 *Suppose that a regular generating function Φ satisfies condition (**iii**):*

$$\mathrm{Re}\, i^n \Theta_{b,\beta}(x,y) = 0 \text{ for any } x, y \in X,\ x \neq y.$$

Then an arbitrary function $f \in L_{2\mathrm{comp}}(X)$ can be reconstructed from data of $\mathrm{M}_b f$ by

$$D_{b,\beta} f = \mathrm{N}_\beta \mathrm{M}_b f + \Theta_{b,\beta} f \tag{10}$$

where $\Theta_{b,\beta}$ is an operator with the kernel $\Theta_{b,\beta}(x,y)$ and for even n,

$$\mathrm{N}_\beta g(x) = \frac{(n-1)!}{\mathrm{j}^n}\int_{Z(x)} \frac{g(\sigma)\beta(x,\sigma)}{\Phi(x,\sigma)^n}\frac{\mathrm{d}\Sigma}{\mathrm{d}_\sigma\Phi},$$

for odd n,

$$\mathrm{N}_\beta g(x) = \frac{1}{2\mathrm{j}^{n-1}}\int_{Z(x)} \delta^{(n-1)}(\Phi(x,\sigma)) g(\sigma)\beta(x,\sigma)\frac{\mathrm{d}\Sigma}{\mathrm{d}_\sigma\Phi}$$

is a bounded operator $H^{(n-1)/2}_{\mathrm{comp}}(\Sigma) \to L_{2\mathrm{loc}}(X)$; and

$$D_{b,\beta}(x) = \frac{1}{|\mathrm{S}^{n-1}|}\int_{Z(x)} \frac{b(x,\sigma)\beta(x,\sigma)}{|\mathrm{d}_x\Phi(x,\sigma)|^n_{\mathrm{g}}}\frac{\mathrm{d}\Sigma}{\mathrm{d}_\sigma\Phi}. \tag{11}$$

A proof is given in [6] for the case $b = \beta = 1$. The general case can be obtained in the same lines.

References

[1] Cormack, A.M.: The Radon transform on a family of curves in the plane I. Proc. Am. Math. Soc. **83**, 325-330 (1981)

[2] Cormack, A.M.: The Radon transform on a family of curves in the plane II. Proc. Am. Math. Soc. **86**, 293-298 (1982)

[3] Cormack, A.M.: Radon's problem - old and new. SIAM-AMS Proceedings, Amer. Math. Soc., Providence **14** 33-39 (1984)

[4] Cormack, A.M.: Radon's problem for some surfaces in $\mathbb{R}^n$. Proc. Am. Math. Soc. **99**, 305-312 (1987)

[5] Cormack, A.M.: A paraboloidal Radon transform. 75 years of Radon transform (Vienna, 1992), 105–109, Conf. Proc. Lecture Notes Math. Phys., IV, Int. Press, Cambridge, MA (1994)

[6] Palamodov, V. P.: A uniform reconstruction formula in integral geometry. Inverse Probl. **28,** 065014 (2012)